EL DORADO COUNTY EMS AGENCY

PREHOSPITAL PROTOCOLS

**EMS Agency Medical Director** 

Effective: January 1, 2017

## BLS Altered Level of Consciousness

## NEEDED INFORMATION:

- Surroundings:
  - Syringes, blood glucose monitoring supplies, insulin
- Change in mental status:
  - Baseline status, onset and progression of altered state, symptoms prior to altered state such as headache, seizures, confusion, and trauma
- Medical History:
  - Diabetes, epilepsy, substance abuse, mental health, medications, and allergies.
- Consider: Stroke and overdose/intoxication

## TREATMENT:

- 1. Reassure patient and place in position of comfort or supine if hypotensive.
- 2. ABC's
- 3. Ensure ALS Response
- 4. Oxygen 10-15 L/min via non-rebreather mask. Patients with ineffective respirations support with BVM ventilation.
- 5. Suction as needed
- 6. Routine Medical Care
- 7. For patients with signs and symptoms of hypoglycemia with the ability to maintain their own airway and swallow without difficulty administer 1 tube of oral glucose paste (15grams of prepared oral dextrose solution). May repeat in 10 minutes or encourage drinking/ eating a sugar containing beverage or food. For patients with their own glucose testing equipment a blood sugar value of less than 70 mg/dl should receive oral glucose.
- 8. For patients experiencing a behavioral emergency treat patient in a calm and reassuring manner using padded or soft leather restraints only to prevent from harming self or others, refer to **Restraint Policy**.